

SPECIAL STANDING ORDERS

This memorandum provides the authority for the Emergency Medical Technicians - Paramedics (EMT-Ps) and Registered Nurses (RNs) employed by or providing volunteer services for the following organizations:

Ashland Fire & Rescue
Mercy Flights

to use these specific written standing orders:

Rapid Sequence Intubation	October 5, 2009
Rocuronium	October 5, 2009 in lieu of Vecuronium.

These specific standing orders shall be in effect January 4, 2010 through June 30, 2010 unless revised or amended. All other standing orders effective July 1, 2009 to July 1, 2010 remain in effect.



Paul S. Rostykus, MD, MPH
Supervising Physician

RAPID SEQUENCE INTUBATION

EMT - P

INDICATIONS:

The preferred method to provide endotracheal intubation after inducing unconsciousness and motor paralysis with medications.

PRECAUTIONS:

Must have an alternate method of airway management available. Succinylcholine chloride may cause malignant hyperthermia or fatal hyperkalemia. Paralysis does not stop the brain's seizure activity. This is a two person procedure.

PROCEDURE:

- **Preparation**
 - IV, cardiac monitor and SpO₂ monitor.
 - Suction.
 - Laryngoscope, ET tubes (2 sizes), stylet.
 - Medications drawn up and labeled
 - Alternate airways – BVM, King Supraglottic Airway, dual lumen airway, percutaneous cricothyrotomy.
- **Preoxygenation** – Hi flow oxygen with non-rebreather mask or bag-valve-mask to maximize SpO₂ – 3 minutes or 8 full breaths
- **Premedication** (optional depending on urgency)
 - Lidocaine 1.5 mg/kg IV push - administered 3 minutes before paralysis if increased intracranial pressure or bronchospasm.
- **Sedation & Paralysis**
 - Etomidate 0.3 mg/kg IV push (0.15 – 0.2 mg/kg IV if elderly, debilitated or hypotensive)
 - Succinylcholine 2 mg/kg (preferred) or Rocuronium 1 mg/kg IV push
- **Protection and positioning**
 - Sellick's maneuver until endotracheal tube placed, confirmed & secured.
 - Patient's head in sniffing position
 - BVM ventilation only if SpO₂ < 90%.
- **Placement and proof**
 - Inflate balloon & secure tube.
 - Auscultation
 - End tidal CO₂ continuous capnography
- **Postintubation management**
 - Midazolam for sedation
 - Maintenance paralysis with rocuronium if adequate sedation not obtainable.

ROCURONIUM (OPTIONAL)

TRADE NAME:

Zemuron

ACTION:

Non-depolarizing skeletal muscle relaxant.

INDICATIONS:

- To provide paralysis (paralyzing dose) for rapid sequence intubation when succinylcholine is contraindicated or if prolonged transport time is anticipated.
- To relieve isolated masseter muscle spasm due by succinylcholine.
- To maintain paralysis after intubation after adequate sedation has been provided.

CONTRAINDICATIONS:

N Known sensitivity to rocuronium.

SIDE EFFECTS & PRECAUTIONS:

Rocuronium causes paralysis, not analgesia or amnesia; all patients must receive adequate sedation before paralysis. Patient will require airway management and ventilation.

ROUTE & DOSAGE:

EMT - P:

- Paralyzing Dose:*** 1 mg/kg IV or IO. Usual adult dose is 70-100 mg.
- Maintenance Dose:*** 0.1-0.2 mg/kg bolus IV or IO bolus as paralysis wears off if sedation with midazolam is not adequate.